

Ready to OWN YOUR LOOK?

Ask your aesthetic
specialist about the
BOTOX[®] Cosmetic
SAVINGS EXPERIENCE

See details on other side of this card



Your
OWN YOUR LOOK
BOTOX[®]
COSMETIC
onabotulinumtoxinA
injection
THERE'S ONLY ONE BOTOX[®] COSMETIC

Actual patient. Results may vary.

IMPORTANT SAFETY INFORMATION (continued)

BOTOX[®] Cosmetic dosing units are not the same as, or comparable to, any other botulinum toxin product.

There has not been a confirmed serious case of spread of toxin effect when BOTOX[®] Cosmetic has been used at the recommended dose to treat frown lines, crow's feet lines, and/or forehead lines.

BOTOX[®] Cosmetic may cause loss of strength or general muscle weakness, vision problems, or dizziness within hours to weeks of taking BOTOX[®] Cosmetic. **If this happens, do not drive a car, operate machinery, or do other dangerous activities.**

Serious and/or immediate allergic reactions have been reported. They include: itching, rash, red itchy welts, wheezing, asthma symptoms, or dizziness or feeling faint. Get medical help right away if you are wheezing or have asthma symptoms, or if you become dizzy or faint.

Do not receive BOTOX[®] Cosmetic if you: are allergic to any of the ingredients in BOTOX[®] Cosmetic (see Medication Guide for ingredients); had an allergic reaction to any other botulinum toxin product such as *Myobloc[®]* (rimabotulinumtoxinB), *Dysport[®]* (abobotulinumtoxinA), or *Xeomin[®]* (incobotulinumtoxinA); have a skin infection at the planned injection site.

Tell your doctor about all your muscle or nerve conditions, such as ALS or Lou Gehrig's disease, myasthenia gravis, or Lambert-Eaton syndrome, as you may be at increased risk of serious side effects including difficulty swallowing and difficulty breathing from typical doses of BOTOX[®] Cosmetic.

Tell your doctor about all your medical conditions, including: plans to have surgery; had surgery on your face; have trouble raising your eyebrows; drooping eyelids; any other abnormal facial change; are pregnant or plan to become pregnant (it is not known if BOTOX[®] Cosmetic can harm your unborn baby); are breast-feeding or plan to (it is not known if BOTOX[®] Cosmetic passes into breast milk).

Tell your doctor about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Using BOTOX[®] Cosmetic with certain other medicines may cause serious side effects. **Do not start any new medicines until you have told your doctor that you have received BOTOX[®] Cosmetic in the past.**

Tell your doctor if you have received any other botulinum toxin product in the last 4 months; have received injections of botulinum toxin such as *Myobloc[®]*, *Dysport[®]*, or *Xeomin[®]* in the past (tell your doctor exactly which product you received); have recently received an antibiotic by injection; take muscle relaxants; take an allergy or cold medicine; take a sleep medicine; take aspirin-like products or blood thinners.

Other side effects of BOTOX[®] Cosmetic include: dry mouth; discomfort or pain at the injection site; tiredness; headache; neck pain; and eye problems: double vision, blurred vision, decreased eyesight, drooping eyelids and eyebrows, swelling of your eyelids and dry eyes.

For more information refer to the Medication Guide or talk with your doctor.

To report a side effect, please call Allergan at 1-800-678-1605.

Please see accompanying Summary of Information about BOTOX[®] Cosmetic.

ENJOY 4 REWARDS WITH BOTOX[®] COSMETIC (onabotulinumtoxinA)

4
Your

OWN YOUR LOOK

BOTOX[®]
COSMETIC
onabotulinumtoxinA
injection

THERE'S ONLY ONE BOTOX[®] COSMETIC

1 + 2
SAVE
\$100*

3
SAVE \$100*
ON JUVÉDERM[®]
3RD
TREATMENT

If you are an appropriate candidate for treatment, you can save \$100 on 2 JUVÉDERM[®] syringes at your third BOTOX[®] Cosmetic treatment.[†]

Rebate may be applied to any 1.0 mL JUVÉDERM[®] product purchase. Treatments must be administered in the same visit. Please note that the safety and efficacy of these products for combined use have not been studied.

UNLOCK ONYX

You'll also be welcomed to an unpublished *Brilliant Distinctions*[®] tier named Onyx. Members get access to:

- Premier savings promotions, local events, and customized partner offers

3
4
COMPLIMENTARY
BOTOX[®] COSMETIC*
4TH
TREATMENT

You can receive your 4th treatment **FREE** when entire series is completed. End date varies based on first treatment start date.

*You must receive each treatment in order. Max of 50 units free. Units over 50 may be charged to you. Injection fee may be added.

[†]Rebate is valid for *Brilliant Distinctions*[®] members only. You must be a *Brilliant Distinctions*[®] member at time of treatment to redeem. Only 1 rebate per member. Rebates cannot be combined with other rebates within the JUVÉDERM[®] Collection, but can be combined with other *Brilliant Distinctions*[®] coupons and rebates. Allergan reserves the right to alter or cancel the program with no advance notice or obligation.

1ST
TREATMENT + 2ND
TREATMENT
\$50 OFF \$50 OFF

After you receive your first treatment, your second rebate is sent directly to your *Brilliant Distinctions*[®] Rewards Program account.

Approved Uses

BOTOX[®] Cosmetic is a prescription medicine that is injected into muscles and used to temporarily improve the look of moderate to severe forehead lines, crow's feet lines, and frown lines between the eyebrows in adults.

IMPORTANT SAFETY INFORMATION

BOTOX[®] Cosmetic may cause serious side effects that can be life threatening. Get medical help right away if you have any of these problems any time (hours to weeks) after injection of BOTOX[®] Cosmetic:

- Problems swallowing, speaking, or breathing, due to weakening of associated muscles, can be severe

and result in loss of life. You are at the highest risk if these problems are pre-existing before injection. Swallowing problems may last for several months.

- **Spread of toxin effects.** The effect of botulinum toxin may affect areas away from the injection site and cause serious symptoms including: loss of strength and all-over muscle weakness, double vision, blurred vision and drooping eyelids, hoarseness or change or loss of voice, trouble saying words clearly, loss of bladder control, trouble breathing, and trouble swallowing.

See additional Important Safety Information on reverse side.